

<https://helda.helsinki.fi>

---

## UV-screening and springtime recovery of photosynthetic capacity in leaves of *Vaccinium vitis-idaea* above and below the snow pack

Solanki, Twinkle

2019-01

---

Solanki , T , Aphalo , P J , Neimane , S , Hartikainen , S M , Pieristè , M , Shapiguzov , A , Porcar Castell , J A , Atherton , J M , Heikkilä , A & Robson , T M 2019 , ' UV-screening and springtime recovery of photosynthetic capacity in leaves of *Vaccinium vitis-idaea* above and below the snow pack ' , Plant Physiology and Biochemistry , vol. 134 , pp. 40-52 . <https://doi.org/10.1016/j.plaphy.2018.09.003>

---

<http://hdl.handle.net/10138/319053>

<https://doi.org/10.1016/j.plaphy.2018.09.003>

---

cc\_by\_nc

submittedVersion

---

*Downloaded from Helda, University of Helsinki institutional repository.*

*This is an electronic reprint of the original article.*

*This reprint may differ from the original in pagination and typographic detail.*

*Please cite the original version.*

Title: UV-screening and springtime recovery of photosynthetic capacity in leaves of *Vaccinium vitis-idaea* above and below the snow pack

Authors: Twinkle Solanki<sup>1</sup>, Pedro J Aphalo<sup>1</sup>, Santa Neimane<sup>1,2</sup>, Saara M Hartikainen<sup>1</sup>, Marta Pieristè<sup>1,3</sup>, Alexey Shapiguzov<sup>1</sup>, J Albert Porcar-Castell<sup>4</sup>, Jon Atherton<sup>4</sup>, Anu Heikkilä<sup>5</sup>, T. Matthew Robson<sup>1</sup>✉

Author Affiliations: <sup>1</sup> Organismal and Evolutionary Biology, Viikki Plant Science Centre (ViPS), Faculty of Biological and Environmental Science, 00014, University of Helsinki, Finland✉

<sup>2</sup> Dept. Plant Physiology, University of Latvia, Jelgavas street 1, LV-1004, Riga, Latvia

<sup>3</sup> Normandie Université, UNIROUEN, Ecodiv URA/EA1293, IRSTEA, FR Scale CNRS 3730, Rouen, France

<sup>4</sup> Viikki Plant Science Centre (ViPS), Department of Forest Sciences, P.O. Box 27, 00014, University of Helsinki, Finland.

<sup>5</sup> Finnish Meteorological Institute (FMI), POB 503, 00101 Helsinki, Finland

Revision for submission to: Plant Physiology and Biochemistry (Special Issue)

**Declaration of Authorship:** TMR SMH conceived & designed the experiments. TS, SN, SMH, MP TMR, did the majority of the field work and sampling, TS, MP, AS, PJA TMR analysed the data, JAPC JA provided site-specific help, and access to meteorological data integral to the experiment, TS, PJA, TMR wrote the manuscript, and all authors provided editorial input.

Abstract (limit 250/250 Words)

Evergreen plants in boreal biomes undergo seasonal hardening and dehardening adjusting their photosynthetic capacity and photoprotection; acclimating to seasonal changes in temperature and irradiance. Leaf epidermal ultraviolet (UV)-screening by flavonols responds to solar radiation, perceived in part through increased ultraviolet-B(UV-B) radiation, and is a candidate trait to provide cross-photoprotection. At Hyytiälä Forestry Station, central Finland, we examined whether the accumulation of flavonols was higher in leaves of *Vaccinium vitis-idaea* L. growing above the snowpack compared with those below the snowpack. We found that leaves exposed to colder temperatures and higher solar radiation towards the top of hummocks suffered greater photoinhibition than those at the base of hummocks. Epidermal UV-screening was highest in upper-hummock leaves, particularly during winter when lower leaves were beneath the snowpack. There was also a negative relationship between indices of flavonols and anthocyanins across all leaves suggesting fine-tuning of flavonoid composition for screening vs. antioxidant activity in response to temperature and irradiance. However, the positive correlation between the maximum quantum yield of photosystem II photochemistry ( $F_v/F_m$ ) and flavonol accumulation in upper hummock leaves did not confer on them any greater cross-protection than would be expected, from the relationship of  $F_v/F_m$  with temperature and irradiance during dehardening present among leaves throughout the hummocks. Irrespective of timing of snow-melt, photosynthesis fully recovered in all leaves, suggesting that *V. vitis-idaea* has the potential to exploit the continuing trend for longer growing seasons in central Finland without incurring significant impairment from reduced duration of snow cover.

41 **Graphical Abstract:** Schematic representing the changes in leaf level processes moderated by the  
42 snowpack, sunlight and temperature after different heights on hummocks of *V. vitis-idaea* during  
43 spring dehardening

44

45 **Keywords** (7 max): spring dehardening; flavonoids; anthocyanins; understory; spectral quality;  
46 photoprotection; Arctic browning

47

## 1. Introduction

An expected outcome of ongoing human-induced climate change is a shift in seasonal precipitation and temperature affecting growing season length (Harsch and HilleRisLambers 2016; IPCC 2014). This has contributed to a reduction in annual snow-cover in the Arctic by as much as 5 days each decade since 1979 (Derksen et al. 2016). At high northern latitudes, less of the winter precipitation is expected to fall as snow and more as rain (Mellander et al. 2007; Rasmus et al. 2004) and temperatures favourable for growth are forecast to occur earlier in the spring (Henry 2008; Jylhä et al. 2008). In Finland, mean annual temperature is expected to increase by 4-11°C and precipitation (and hence cloudiness) by 10-60 % by the end of 21<sup>st</sup> century, with most increases occurring during the winter (Finland's Seventh National Communication under the United Nations Framework Convention on Climate Change, 2017). The benefit of this change for plant growth and photosynthesis is uncertain because warmer mean temperatures together with reduced precipitation as snow may lead a smaller snow pack that melts earlier (Kreyling et al. 2012). The absence of snow cover will imply greater temperature fluctuations close to the ground, exposing plants to colder minimum, and warmer maximum, temperatures when they are starting to deharden at the end of winter (Blume-Werry et al. 2016; Preece and Phoenix, 2013).

Plants possess several photoprotective mechanisms which mitigate abiotic stresses under harsh conditions: (1) when reaction centres are open photochemical quenching ( $q_P$ ) entails activation of the enzymes and metabolites involved in charge separation to produce energy via photosynthesis; (2) excess light energy, causing reaction centres to close, is dissipated through xanthophyll-cycle pigments (carotenoids) zeaxanthin and antheraxanthin in a process regulated by the pH the thylakoid lumen (Adams et al. 2004; Murchie and Lawson 2013); (3) additionally antioxidants such as phenolic compounds are accumulated to quench reactive oxygen species (ROS) (Agati et al. 2013; Hideg et al. 2013; Salter et al. 2017). It is reasonable to assume that the pressure on these mechanisms is increased when a plant is subject to both fluctuating temperatures and irradiances above the snowpack rather than the more stable conditions offered under snow (Loik et al. 2004; Preece and

Phoenix, 2013). Low temperatures dampen physiological processes meaning that even moderate irradiances cause photoinhibition through photoinactivation via the  $\text{Mn}_4\text{CaO}_5$  cluster or inhibition of the photosystem II ( $\text{PS}_{\text{II}}$ ) electron acceptors (Neuner et al. 1999; Saarinen et al. 2011). Sunlight can penetrate the snowpack, where temperature fluctuations are small, potentially allowing some photosynthesis at low irradiances (Saarinen et al. 2016). Some studies have found the relative transmittance of different regions of the spectrum by snow to be quite even (Gorton and Vogelmann, 2003) while others have reported spectrally-differential attenuation through the snowpack (Curl Jr et al. 1972; Richardson and Salisbury, 1977). Although solar irradiance, particularly ultraviolet-B (UV-B: 280-315 nm) radiation, is low in winter, the ratio of ultraviolet (UV-B plus UV-A: 280-400 nm) to photosynthetically active radiation (PAR: 400-700 nm) in forest understorey shade is enriched (Jansen et al. 2017; Hartikainen et al. 2018).

Flavonoids such as flavonols and anthocyanins are the most-common class of phenolic compounds produced in response to UV-radiation and blue light (Searles et al. 2001; Sullivan et al. 2007; Taulavuori et al. 2016). Flavonols accumulate in the leaf adaxial epidermis absorbing UV-radiation so screening the photosynthetic apparatus in the mesophyll below (Barnes et al. 2017; Caldwell et al. 1983; Nybakken et al. 2004). Although their absorption spectrum is predominately in the UV region, it has been proposed that flavonoids provide cross-protection to high solar irradiance in general (Jansen et al. 2017), and a positive correlation is sometimes reported between flavonoid accumulation and the yield of  $\text{PS}_{\text{II}}$  (Robson et al. 2014; Wargent et al. 2011). This relationship may be attributable to protection against  $\text{PS}_{\text{II}}$  photodamage or enhanced ROS scavenging conferred by flavonoids (reviewed by Takahashi and Badger, 2011). The accumulation of flavonoids in response to UV-radiation is moderated by temperature, being less pronounced in warm than in cool temperatures (Bilger et al. 2007; Coffey et al. 2017; Petridis et al. 2016) where flavonoids can be important in mitigating against oxidative stress (Lütz et al. 2005) and can increase frost hardiness (Chalker-Scott and Scott 2004; Shulze et al. 2016). Hence, the accumulation of flavonoids could provide a feasible

strategy whereby evergreen plants ameliorate photoinhibition, imposed by increasing irradiance and temperature fluctuations, as they deharden over the spring.

Boreal evergreen and wintergreen plants adjust light harvesting complexes to optimise quenching processes at the expense of the light harvesting and photosynthetic capacity of their leaves during winter when light and temperature conditions are unfavourable (Adams et al. 2004). Subsequently, they undergo a recovery period in preparation for more favourable growth conditions in late spring and summer (Muller et al. 2005). During this period of dehardening, full functionality returns to the photosynthetic apparatus of leaves and this can be measured using chlorophyll fluorescence as an increase in the yield of PS<sub>II</sub> (Murchie and Lawton, 2013; Saarinen et al. 2011). Monitoring leaf-level chlorophyll fluorescence of evergreen trees (Ensminger et al. 2004; Porcar-Castell et al. 2008) and field-layer plants (Starr and Oberbauer, 2003) through spring in the boreal forest, has revealed that recovery of  $F_v/F_m$  is modulated by temperature and light.

Here, we aimed to compare leaf epidermal flavonoids (flavonols and anthocyanins), chlorophyll content and the PS<sub>II</sub> yield at different positions on hummocks relative to the snowpack, through a seasonal time-course of measurements of evergreen-shrub *Vaccinium vitis-idaea* L. During the spring dehardening period, the over-wintering leaves of this species are adapted to take advantage of favourable early-spring conditions by photosynthesizing when still under the snowpack (Lundell et al. 2008; Saarinen et al. 2011; Starr and Oberbauer, 2003). Our hypothesis was that leaves at the top of the hummock above the snowpack during winter and early spring would suffer greater photoinhibition because they encounter fluctuating temperatures and higher irradiance than those at the base of hummocks receiving lower irradiance, of both PAR and UV radiation, and more stable temperatures remaining closer to 0°C under the snow. However, we also expected higher accumulation of epidermal UV-absorbing compounds (flavonols and anthocyanins) in leaves above the snow, and for this to partially ameliorate the photoinhibition they suffered relative to those at equivalent temperatures beneath the snowpack.

## 2. Methods

### 2.1. Study site and species

The study was performed under an open *Pinus sylvestris* L. canopy, basal area 23 m<sup>2</sup>ha<sup>-1</sup>, on level ground at Hyytiälä Forestry Station of the University of Helsinki in central Finland (61.84741°N 24.29477°E 181 m asl). The hummock-forming evergreen shrub *V. vitis-idaea* covers vast areas of peatland and boreal forest at high latitudes in the Northern Hemisphere (Karlsson, 1985; Lundell et al. 2008). At our site, the understorey comprises almost exclusively *V. vitis-idaea* in hummocks of up to 75 cm in height intermingled with *Vaccinium myrtillus* L. Four hummocks, positioned at least a metre from the nearest pine tree, were selected at random within a 10-by-10-m area of understorey. These hummocks were divided vertically into three height categories on 14-02-2017: the “upper” hummock being the uppermost shoots, most exposed to the wind and sun, the “middle” hummock being the outer shoots on the south side of the hummock, somewhat sheltered from the wind but just below the snow pack through the winter, and the “lower” hummock being shoots at the very base of the hummock on the south side, below the snow pack during the whole winter and early spring.

The snow depth relative to the lower, middle and upper hummocks, and the leaf phenology of the *Vaccinium* species, was monitored throughout the study concomitantly with leaf traits, and these data were compared against a continuous monitoring by Phenocam time-lapse images nearby in the stand (Kolari et al. 2017). By 19-06-2017, plants of both *V. vitis-idaea* and *V. myrtillus* produced new leaves. From this date onwards, until our final measurement date, 08-11-2017, a set of measurements was also made on these new *V. vitis-idaea* leaves (unfurling in June 2017) in same way as those on leaves produced in June 2016. This allowed us to obtain data for leaves from the time of their unfurling through their first and second years of growth. To allow this continuum of leaf age to be visualised in the figures we have combined these two set of leaf traits in a single time series; with 2017 leaves plotted from their unfurling until 08-11-2017 (day 312 of the year, leaf age 152 days) and 2016 leaves from 14-02-2017 onwards (day 45 of the year, leaf age 250 days).



## 2.2. Characterisation of the forest canopy by hemispherical photograph

Plant area index (PAI)( $\text{m}^2\text{m}^{-2}$ ), was estimated at each of the four hummocks by taking hemispherical photographs (at 40 cm height from the ground) through a Sigma 4.5mm f2.8 EX DC HSM circular Fisheye lens (Sigma Corporation of America, Ronkonkoma, NY) on a Nikon D7000 camera (Nikon corporation, Tokyo, Japan) levelled on a tripod. Measurements were done on a completely overcast day in mid-July (20-07-2017) to ensure homogeneous grey sky and a good contrast between the sky and the canopy (Fig S1). All photos were analysed following the procedure detailed in Hartikainen et al. (2018).

## 2.3. Measurement of spectral irradiance at the study site

The solar spectral irradiance was measured close to solar noon, directly to the south of each of the four hummocks on 07-03-2017 and 19-06-2017 in sunflecks and canopy shade on completely clear-sky days. These data were compared with corresponding measurements in an open field next to the forest stand earlier on the same days. Spectral irradiance recordings were made using a Maya 2000 Pro (Ocean Optics, Dunedin, FL, USA) array spectrometer. The cosine diffuser (Bentham Instruments Ltd. Reading UK) was levelled on the tripod 40-cm high and attached with a fibre-optic cable to the spectrometer. Measurements were corrected for dark-noise and stray light in the UV range ( $<400\text{ nm}$ ) using a protocol detailed by Hartikainen et al. (2018). The spectrometer had been calibrated against a Bentham scanning spectroradiometer for outdoor measurements of solar spectral irradiance, over the wavelength range 290-900 nm. Pre-processing of the data was performed using functions from the R packages Ooacquire and Photobiology in R version 3.4.3 (R Core Team, 2018).

## 2.4. Optical measurements of *V. vitis-idaea* leaf pigments and photosynthetic capacity

On each measurement date, starting 14-02-2017, then 07-03-2017, and at 2-week intervals subsequently until 08-07-2017, with an additional final measurement date of 08-11-2017, leaf optical properties and photosynthetic capacity were measured. Ten leaves (remaining attached to the plants) were measured *in situ* from each height (upper, middle, lower) on four hummocks. Sun-exposed leaves of equivalent size were measured (or leaves which would have been sun exposed had it not been cloudy on overcast days), all from different shoots of each *V. vitis-idaea* hummock, avoiding the

three distal leaves of a shoot. Lower and middle hummock leaves were, when necessary, excavated from beneath the snow to be measured and subsequently the snow was replaced over these leaves. Leaves that were damp from dew, rain or melting snow were quickly but gently dried with a tissue prior to measurement.

Dualux Scientific<sup>+</sup> (Force-A, University Paris-Sud, Orsy, France) was used to optically assess the epidermal flavonol concentration (epidermal absorption in the UV-A region at 375 nm) and anthocyanin concentration (absorption at 515 nm), from both the adaxial and abaxial sides of each leaf, as well as the leaf chlorophyll concentration (assessed optically by transmittance in the red at 650 nm vs. near-infra red at 710 nm, Cerovic et al. 2012). We recorded the maximum quantum yield of PS<sub>II</sub> photochemistry ( $F_v/F_m$ ; where  $F_v$  [variable fluorescence] is the maximal fluorescence,  $F_m$ , minus the minimal fluorescence,  $F_o$ , of dark-adapted leaves, as defined by Murchie and Lawson, 2013) measured with a mini-PAM (Heinz Walz GmbH, Effeltrich, Germany) following 30 min of dark-adaptation with darkening clips. According to Bokhorst et al. (2010), 30 min in darkness should be sufficient for full dark adaptation of *V. vitis-idaea* leaves year 'round. We also recorded the effective quantum yield of PS<sub>II</sub> photochemistry ( $\phi PS_{II}$  calculated as  $F_q'/F_m'$ ; where  $F_q'$  is the maximal fluorescence,  $F_m'$ , minus the steady-state fluorescence,  $F'$ , of leaves in ambient sunlight, as defined by Murchie and Lawson, 2013) of leaves in the sun and shade in the mid-to-late morning. Where possible, the same ten leaves were used for chlorophyll fluorescence and Dualux measurements on any given day, but it was not feasible to measure the same ten leaves from one measurement date to another, so from day to day leaves meeting the above criteria were selected at random.

## 2.5. Harvest of leaves for trait measurements

On 08-07-2017, leaves of age 29 days and 394 days were harvested. From each age cohort 8-10 leaves were sampled from each of the four hummocks at each of the three heights (2 epidermis × 10 leaves × 3 heights × 4 hummocks). Fresh leaves were placed in zip-lock plastic bags, weighed and digitally photographed against an area reference on the day of sampling, then dried at 40°C for 9 days in a drying oven with forced ventilation before reweighing to obtain their dry weight. Leaf areas were

calculated from the photographs using ImageJ (US National Institute of Health, Bethesda, Maryland, USA: following Wang, 2017) and used to calculate the leaf mass area (LMA  $\text{mg mm}^{-2}$ ). The leaf area covering the optical window of the Dualex Scientific <sup>+</sup> was  $47.7 \text{ mm}^2$  (Area =  $\pi r^2$ ,  $r = 3.90 \text{ mm}$ ), used with LMA, to calculate per area and per mass values of pigment concentrations on the harvest dates.

## 2.6. Environmental monitoring of temperature during the experiment

At the time of each measurement of leaf optical properties, the temperature of the leaves of *V. vitis-idaea* at each height on all four hummocks was measured at close range with a high precision infra-red thermometer with special macro optics (LS laser-sight Optris GmbH, Berlin, Germany). Measurements were also recorded at the tree base and at 1 m 50 cm height on the trunks (emissivity  $c 0.98$ ) of the four nearest trees to the hummocks at opposite compass points (North and South at midday) in the sun and shade, and on the snow surface (emissivity of tree trunks and snow  $c 0.98$ ).

Meteorological data for the experimental period were obtained from the SMEAR II ICOS Hyytiälä station through the AVAA data research portal (<https://avaa.tdata.fi/>). Temperature in the understorey was recorded every minute with a PT100 sensor (Pentronic, Västervik, Sweden) mounted on a mast at 4 m height and soil surface temperature was recorded every minute with a KTY 81-110 sensor (NXP Philips, Amsterdam, Netherlands). Understorey photosynthetically active radiation (PAR) was measured with an array of four Li190SZ PAR sensors (LI-COR, Lincoln, NE) at 0.6m height next to the hummocks. These time series of meteorological data were used to calculate averages over a range of time periods preceding each measurement (1 hour, 1 day, 2 days, and 2 weeks) and the relationship between temperature and PAR over these periods and leaf trait data was plotted. Of these four time integrals the best fit to the trait data was retained.

## 2.7. Data Analysis

Differences due to hummock height among optically-assessed leaf traits were tested using a two-way ANOVA with hummock height, and leaf age (days) as fixed factors, or a three-way-ANOVA for the responses of flavonols and anthocyanins which were measured in both the adaxial and abaxial epidermis of each leaf (with leaf side as a nested third fixed factor). One measurement date (Leaf Age:

250 days), when only the adaxial epidermis was measured with the Dualex, was excluded from the three-way ANOVA. The equivalent single-factor ANOVA was used to test the effect of hummock height on traits measured only once, e.g. from harvested leaves at the end of the experiment. The four hummocks were the experimental units of replication and height was nested within each hummock. When significance differences were produced by the ANOVA, pairwise differences were identified between the responses across three levels of hummock height on either leaf side on each measurement date (function `glht` in the package `Multcomp`). Holm's adjustment was used to account for multiple pairwise comparisons. The relationship of  $F_v/F_m$  with temperature and PAR was assessed using linear regression (function `lm()`). All the analyses were performed using R version 3.4.3 (R core development team, 2018).

### 3. Results

#### 3.1. Forest cover, snow depth and temperature

The snow depth was c. 40 cm at the first winter measurement (14-02-2017), meaning that the upper shoots of hummocks (upper hummocks) were above the snow, the middle of the hummocks were just below the top of the snowpack, and lower shoots of hummocks (lower hummocks) were entirely beneath the snowpack. From 14-02-2017 onwards, the snow depth didn't increase from this maximum and remained above the middle-hummock shoots until 27-03-2017 and above the lower-hummock shoots until snow melt was complete on 10-05-2017.

Soil temperatures under the snow were relatively stable, starting to fluctuate only when snow melt was almost complete and the lower hummock leaves above the snow (Fig. 1; Table 1). From the time series of snow depth and air temperature data, compared with IR-thermometer measurements of snow-pack temperature (Fig. 1), we were able to estimate that the upper-hummock leaves were subject to 111 freeze-thaw cycles, the middle-hummock leaves 86 and lower-hummock leaves 38 such cycles, during the period between New Year and mid-summer 2017 which encompasses full dehardening (Table 1).

### 3.2. Spectral irradiance at the study site

The solar spectral irradiance was measured on three dates during the winter and spring (14-02-2017, 07-03-2017 and 19-06-2017). On all three dates, the normalised spectra show that PAR was depleted in the under-canopy sunflecks and shade compared with the open field, and the relative contribution of UV radiation was similar in the sunflecks and open sun but enriched in the shade (Figs. 2 and S2, Table S1). The spectral composition of the sunflecks varied with sunfleck size as well as time of year, and the relative contribution to shade by needles and branches also affected the shape of these spectra. The Plant Area Index (PAI) and sunfleck duration on 20-07-2017 were  $2.14 \pm 0.05 \text{ m}^2 \text{ m}^{-2}$  and  $169.6 \pm 11.27 \text{ min day}^{-1}$  respectively, and was similar among the four hummocks (Table S2).

The reflectance (up to 90%) and transmittance of sunlight by the snowpack modifies amount and composition of radiation received by leaves above and below the snowpack at different hummocks heights. We found the snowpack to transmit about 10% of incident PAR irradiance to a depth of 10 cm, and a similar or slightly higher percentage of UV radiation to reach the same depth (Robson unpublished data) in line with comparable measurements elsewhere (Richardson and Salisbury, 1977; Gorton and Vogelmann, 2003; Perovich 2007).

### 3.3. Seasonal changes in leaf pigments at different heights on the *V. vitis-idaea* hummocks

Leaf flavonoid concentration (i.e. both flavonols and anthocyanins), measured optically and presented as an index of absorption on a per leaf-area basis, was most similar directly after leaf unfolding both among hummocks and between leaf sides. As the leaves matured, their flavonol concentration increased in the adaxial epidermis and initially decreased in the abaxial epidermis between the first and second measurement of new leaves during the summer (Leaf Age 10-29 days, Fig. 3A). During this period, UV-screening by flavonols was greater in the upper-, and middle-, hummock leaves than the lower leaves. Accumulation of flavonols continued until the start of winter in the adaxial-, and to a lesser extent the abaxial-, epidermis, and only started to definitely decline with age following the spring and summer of their second year (Leaf Age 271 days; Fig. 3A). Leaves of

*V. vitis-idaea* can remain productive for 5 years or more, but after their first growing season maximum photosynthesis declines sharply (Karlsson, 1985).

Adaxial epidermal flavonol concentration in *V. vitis-idaea* leaves transiently decreased during winter in lower- and mid-hummock leaves when they were beneath the snowpack, but started to increase again from 24-03-2017 (Leaf Age 288 days) coinciding with temperatures rising above freezing. In sequence, the upper, then middle, then lower hummock leaves' adaxial flavonol concentration increased around the time that the middle hummocks leaves emerged from the snowpack (Fig. 3A, from 27-03-2017, Leaf Age 291 days). This phase of flavonol accumulation continued until the end of the snow melt period (16-05-2017 Leaf Age 341 days), after which flavonol concentration was stable and eventually started to decline from 19-06-2017 Leaf Age 375 days (Fig. 3A). Flavonol accumulation in the adaxial epidermis during spring was greater than that in the abaxial epidermis, leading the adaxial-to-abaxial flavonol ratio to increase until 08-07-2017, Leaf Age 394 days, but it returned to the equivalent of pre-spring ratios later in the year (08-11-2017, Leaf Age 517; Figs. 3A and S3A).

Leaf anthocyanin concentration is typically highest in young leaves, and higher in the abaxial than the adaxial epidermis, declining once leaves are mature (Landi et al. 2015): our results were consistent with this pattern until late winter, when anthocyanin accumulation rebounded (07-03-2017, Leaf Age 271 days, Fig. 3B). Initially, there were no differences in anthocyanin concentration with respect to height on the hummocks, and this remained the case until 24-03-2017 (Leaf Age 313 days) after the middle-hummock had emerged but while the lower hummock was still beneath the snow pack. During this period (24-03-2017 to 18-04-2017), adaxial epidermal anthocyanins continued to accumulate in the upper hummock leaves (Fig. 3B) and the difference in adaxial epidermal anthocyanin concentration with height was maintained, with higher values in upper hummock leaves compared with the others (Fig. 3B). In the abaxial epidermis, a corresponding difference with height occurred later during the 6-week spring dehardening period (19-04-2017 to 30-05-2017: Leaf Age 313-

355 days), persisting until abaxial anthocyanin concentration was maximal in the upper and mid hummock leaves between (17/30-05-2017, Leaf Age 341-355 days). By this time, the abaxial epidermal anthocyanins had already started to decline in the lower-hummock (Fig. 3B). By mid-summer, both adaxial and abaxial epidermal anthocyanins had converged to similar values across the three height classes (19-06-2017: Leaf Age 375) (Fig. 3B). The ratios of adaxial-to-abaxial epidermal anthocyanins all declined throughout the spring, but differed among the three heights, being highest in the upper-hummock leaves and lowest in the lower-hummock leaves (Fig. S3B).

Chlorophyll concentration per leaf area increased in young leaves until 14-02-2017 (Leaf Age 250 days), by which time it was lower in the lower-hummock leaves than the middle-and-upper-hummock leaves (Fig. 3C). This difference persisted until the upper and middle hummocks were above the snowpack but the lower hummock beneath the snowpack (18-04-2017, Leaf Age 313 days), at which time chlorophyll concentration of the lower-hummock leaves increased beyond that of the exposed upper and middle hummock leaves where it declined (Fig. 3C). Lower-hummock leaves retained the highest chlorophyll concentration from then until midsummer by which time chlorophyll values at the three heights converged again (19-06-2017, Leaf Age 394 days) (Fig. 3C).

The ratio leaf flavonols vs. anthocyanins (sum of both epidermis) was higher in the upper hummocks than the lower hummocks through the winter (Fig. 4A). In the adaxial epidermis, the rate of flavonol accumulation was faster than that of anthocyanins during the dehardening period so this ratio rose, particularly in the lower hummock leaves (due to increasing flavonol accumulation as described above - Fig 4A). There was negative relationship between flavonols and anthocyanins in the adaxial epidermis when data are plotted for all leaves and all dates (Fig. 4B). This relationship may be confounded by leaf age but nevertheless persisted for the adaxial epidermis when the relationships were calculated separately for each measurement date (Table S5).

### 3.4. Seasonal changes in photosynthetic yield of PS<sub>II</sub> in leaves at different heights on the *V.*

#### *vitis-idaea* hummocks

The  $F_v/F_m$  of leaves across the three heights on the hummocks attained their maximal values of about 0.82 on reaching maturity on their second measurement following unfurling (09-07-2017, Leaf Age 30 days) and  $F_v/F_m$  remained high prior to snow cover in the winter (Fig. 5A). During the winter and spring,  $F_v/F_m$  was depressed most in the upper hummocks, then middle hummocks and least depressed in the lower hummocks (Fig. 5A). This ranking of  $F_v/F_m$  according to height on the hummocks was retained throughout the spring dehardening period (until 30-05-2017: Leaf Age 355 days), but by midsummer one-year old leaves at all three heights had attained similar maximal  $F_v/F_m$  again (19-06-2017, Leaf Age 375 days) (Fig. 5A).

The  $\phi PS_{II}$ , measured *in situ* under ambient sunlight, varied with the solar irradiance, i.e. on clear-sky measurement days  $\phi PS_{II}$  was very low during the dehardening period (Fig. 5B). At sub-zero temperatures (e.g. -5.5 °C on 07-03-2017, Leaf Age 313 days) with clear-sky irradiance during early spring, the  $\phi PS_{II}$  was lowest and approached zero. Though bear in mind that it was necessary to remove the snow from covered leaves during winter to take the measurement, exposing them briefly to higher PAR than they would otherwise have received and thus probably caused greater photoinhibition than when under the snow. Nevertheless, in early spring on days 83 and 96 (leaf age 288 and 301 days),  $\phi PS_{II}$  of almost 0.2 was reached in the lower hummock leaves (Fig. 5B). Both measures of photosynthetic capacity ( $F_v/F_m$  and  $\phi PS_{II}$ ) responded to day-to-day differences in environmental conditions, as evident from differences between  $F_v/F_m$  values from consecutive daily measurements made on three occasions during the spring dehardening period (Fig. 5B, Leaf Age 313-314, 327-328, 341-342 days).

### 3.5. The relationship between *V. vitis-idaea* photosynthetic yield, leaf flavonoids and environmental conditions

We tested the relationship between temperature and  $F_v/F_m$  for leaves from the upper, middle, and lower hummocks, identifying their position relative to the snowpack prior to each  $F_v/F_m$



measurement. The air temperature ( $T_{aw}$ ) or soil temperature ( $T_{sw}$ ) at the site were positively correlated with the average  $F_v/F_m$  of leaves above or below the snowpack respectively, and in order to make a direct comparison with other studies of *V. vitis-idaea* (Bokhorst et al. 2010; Saarinen et al. 2016), we tested a range of different time periods to find the closest relationship. Of these time periods, the temperature over the 2 weeks prior to each  $F_v/F_m$  measurement gave the best fit; for leaves above the snowpack vs.  $T_{aw}$  (adj.  $R^2=0.91$ , Table S4), and leaves below the snowpack vs.  $T_{sw}$  (adj.  $R^2 = 0.81$ , Table S4). These relationships with temperature were stronger than  $F_v/F_m$  vs. PAR on the day of measurement for leaves above the snowpack (adj.  $R^2 = 0.74$ , Table S4).

Individual leaf measurements of  $F_v/F_m$  were also positively correlated with leaf temperature (Fig. 6A) and PAR incident on the leaf (Fig. 6B) at the time of measurements, and likewise with adaxial-epidermal flavonol concentration on the day of measurement (Fig. 6C). However, these relationships were strongly conditioned by measurement day and held only during the period from 24-03-2017 to 17-05-2017: Leaf Age 281-342 days (Fig. 6). The relationships broke down once spring recovery of  $F_v/F_m$  was almost complete (30-05-2017 Leaf Age 355 days – not shown) and in winter when leaf temperatures were consistently around 0°C. During dehardening, all three relationships were strongest in the upper hummocks and remained strong in the middle hummocks, where  $F_v/F_m$  recovery coincided with an increase in adaxial epidermal flavonols during this time period, but were but much less strong in the lower hummocks (Fig. 6C). The same relationship was not evident for anthocyanins in either epidermis (not shown).

## 4. Discussion

### 4.1 Consistent pattern of increasing flavonoids with hummock height throughout the year

Differences in exposure to harsh weather conditions among adjacent plants or shoots may be particularly pronounced in the understorey of open canopies typical of the boreal forest which unevenly intercept solar radiation (both incoming and reflected or emitted radiation) and snow (Hedstrom and Pomeroy, 1998). The undulating topography created by hummock-forming plants, adds to the heterogeneity of irradiance, temperature, and snow cover (Essery et al. 2008; Pomeroy et

al. 2002; Sturm et al. 2001). Accordingly, a large proportion of their leaves can remain above the snowpack over winter where they are subject to greater fluctuations in temperature and irradiance than those beneath the snowpack. In our study, this was the case for the upper leaves of *V. vitis-idaea* hummocks, which accumulated more epidermal UV-screening (attributable to flavonols) than lower hummock leaves. Using a Dualex to optically detect UV-screening allowed us to repeatedly and non-destructively follow changes over time on the same plant but not to gather biochemical information on flavonoid composition.

The results of studies based on optical properties of the leaf epidermis *in vivo* and measurements of whole-leaf extracts can be difficult to reconcile because the two methods are not equivalent. In *V. vitis-idaea*, flavonoids can be found in the trichomes and bound to the epidermal cell walls (Semerdjieva et al. 2003a, 2003b) where they will affect UV-screening, whereas they are spread throughout the leaf including chloroplasts in the mesophyll and cuticular waxes in species like *V. myrillus* (Semerdjieva et al. 2003b). Despite a certain lack of equivalency, it is worth noting that the relative differences we found in UV-screening within *V. vitis-idaea* hummocks, among leaves differing only in the time spent above and below the snowpack (Fig. 1), were larger than those produced in the same species by a 25% UV supplementation (Semerdjieva et al. 2003b). A strong negative correlation between accumulation of leaf flavonoids and increasing temperature can obscure the effects of UV-B radiation (Coffey et al. 2017), and consequently, the differences in temperature across hummock height related to differences in seasonal snow cover may partially explain the vertical gradient in flavonols accumulation we report. It has not yet been elucidated whether both temperature and UV-B radiation may regulate flavonoid accumulation via the *uvr8* pathway, but this would constitute an interesting future avenue of research.

The response of *V. vitis-idaea* leaves to UV radiation at high latitudes has already been extensively studied in the context of northern-hemisphere stratospheric ozone depletion in the late 1990s and 2000s (e.g. Johansson et al. 1995; Phoenix et al. 2001; Semerdjieva et al. 2003a). The

leaves of evergreen species emerge from snow cover in spring when seasonal UV-B exposure can be periodically high, but a meta-analysis generally found evergreen species like *V. vitis-idaea* to be resilient against such seasonal increases in UV radiation (Newsham and Robinson, 2009). The modest flavonoid induction caused by increased UV-B radiation could help protect from photodamage to PS<sub>II</sub> and improve antioxidant capacity during dehardening at the start of the growing season when high irradiance and cold temperatures in unison exacerbate photoinhibition (Takahashi and Badger 2011).

Most studies from the 1990s-early-2000s have presented these data on a per mass basis which is not as easily interpretable as leaf-area based measures in terms of screening. Differences in LMA among the one-year old leaves in our experiment caused the effect of hummock height on flavonols and anthocyanins per-leaf-area to be reduced when considered on a per-mass basis (Table 2). Likewise, the effect of UV-B radiation on flavonol accumulation is sometimes smaller on a per-leaf-mass than per-leaf-area basis in understorey shrubs, because UV-B exposure causes mature leaves to be thicker and tougher (Newsham and Robinson, 2009, Semerdjieva et al. 2003a).

The increase in the ratio of adaxial to abaxial epidermal flavonols in the lower hummock leaves (Fig. S3A) during snow melt suggests that accumulation of adaxial flavonols could aid in photoprotection in addition to amelioration of cold temperature stress as antioxidants (Chalker-Scott and Scott, 2004). Comparison of the UV screening on either side of a leaf show this relationship to be highly species specific and environmentally dependent: sometimes UV screening by the adaxial and abaxial epidermises are strongly positively correlated (Bartlöd et al. 2007, Siipola et al. 2014) sometimes this relationship is weak (Kolb and Pfündel 2005, Morales et al. 2011), as was the case in our study. The accumulation of epidermal flavonoids is commonly attributed to exposure to UV-B radiation, or high irradiance of UV-A radiation and blue light (Barnes et al., 2017), although flavonoid accumulation is also known to increase in response to cold temperatures (Agati et al, 2013). Temporal differences in flavonoid (both flavonol and anthocyanin) accumulation and their

localization in the planophile leaves of *V. vitis-idaea* might reflect differences in the allocation of resources based on differences between the adaxial and abaxial epidermis in their relative exposure to cold and high irradiance.

Over the whole dataset, there was an inverse correlation between the anthocyanin and flavonol concentrations in the adaxial epidermis, even though during spring dehardening both tended to increase (Fig. 4). A similar pattern showing a trade-off between these two phenolic compounds was reported in *Arabidopsis thaliana* under controlled conditions, where the ratio of flavonoids to anthocyanins was higher when they received either blue light or UV-A radiation (Brelsford et al, 2018). Anthocyanins are typically found in greater amounts in young leaves and have a specific role as antioxidants involved in cold and high-light tolerance (Havaux and Kloppstech, 2001; Landi et al. 2015). Although anthocyanins only infrequently respond to elevated UV-B radiation (Searles et al. 2001), both flavonols and anthocyanins are phenolic compounds of the flavonoid pathway and it is possible that regulation of this pathway shifts during spring to emphasize the photoprotective function of flavonols driven by seasonally-increasing irradiance of blue light and UV radiation at the expense of anthocyanins whose antioxidant function would be more important during winter (Agati et al. 2013).

#### 4.2 Prolonged snow cover accelerates recovery of photosynthetic capacity during spring dehardening

The timing of spring dehardening in our study was better correlated with temperature than irradiance, and the relationship with temperature was maintained over 2 weeks prior to measurement (Fig 6), in agreement with Karlsson (1985). Several studies have found a consistent relationship between the ambient temperature and  $F_v/F_m$  of *V. vitis-idaea* leaves across a range of latitudes (Bokhorst et al. 2010; Saarinen et al. 2016; Table S6 and S7) and in *Pinus sylvestris* at our Hyytiälä site (Porcar-Castell et al. 2011). Saarinen et al. (2016) report that  $F_v/F_m$  in leaves of *V. vitis-idaea* plants was depressed by snow removal or even when covered by only a thin layer of snow. In our study, upper-hummock leaves had the lowest  $F_v/F_m$  and mid-hummock leaves, which were also only thinly

snow-covered, had lower  $F_v/F_m$  than lower-hummock leaves. Prior to their emergence from under the snow, the  $F_v/F_m$  of lower-hummock leaves was already increasing. It is conceivable this recovery was aided by the low to moderate irradiance that penetrates the upper cm's of the snowpack, potentially activating photoprotective mechanisms during snowmelt (Richardson and Salisbury, 1977). This could partially explain why the difference in  $F_v/F_m$  was maintained between leaves at different heights on the hummocks for several weeks following snow melt rather than the lower leaves exhibiting a cold-shock response upon exposure.

Large fluctuations in temperature and sustained winter warming can both prematurely activate dehardening creating the potential to bring forward the growing season but also increasing the risk of damage if cold winter temperatures return (Bokhorst et al. 2008). Photoinhibition of exposed shoots was high in our study, and leaf chlorophyll concentration dropped in the upper and middle hummock leaves during the spring dehardening period but remained stable in the lower hummock leaves (surpassing that in the upper hummocks) following emergence from under the snowpack. The regulation of leaf chlorophyll may be a means of acclimation, to reduce photon capture under high irradiance to balance the limited carboxylation capacity at low temperature in the upper and middle hummocks when photoprotective mechanisms were surpassed.

Winter- and spring-time depression of  $F_v/F_m$  and  $\phi PS_{II}$  was largely attributable to a decrease in maximal fluorescence ( $F_m$  and  $F_m'$  respectively) rather than an increase in  $F_0$  or  $F_0'$  (Fig. S5). This suggests that the reduction in photosynthetic capacity is the result of decreased light harvesting by  $PS_{II}$  lowering the  $F_m$  and  $F_m'$ . Such a result has previously been attributed to smaller antenna size or sustained NPQ associated with decreased epoxidation of xanthophylls or  $PS_{II}$  reaction centre quenching under cold temperatures (Ivanov et al. 2008), rather than damage or disassembly of the  $PS_{II}$  LHC<sub>II</sub> core subunit super-complexes which would be reflected in higher  $F_0$  or  $F_0'$  (Öquist and Huner, 2003). The positive correlation between epidermal flavonols and  $F_v/F_m$  found in the upper and middle, but not lower, hummock leaves during dehardening suggests that these leaves are acclimating to the

more-exposed conditions. This is consistent with some mechanistic studies which report flavonoids to determine of cold acclimation and frost tolerance of *Arabidopsis thaliana* (Havaux and Klopstech, 2001; Shulz et al. 2016). However, knowledge of the contribution of xanthophyll cycle pigments at different hummock heights, which was not quantified in this study, would allow us to better describe this response (Adams et al. 2004).

#### 4.3 Ecological consequences of changing snow cover for *V. vitis-idaea*

In our study in central Finland, all leaves achieved full functionality following dehardening. Likewise, Saarinen et al. (2016), found that exposed leaves of *V. vitis-idaea*, subjected to experimental removal of the snow pack during winter, fully recovered  $PS_{II}$  capacity during the spring in southern Finland, abet later than those from which snow was not removed. This result is consistent with other Arctic shrub species, e.g. *Rhododendron ferrugineum* L, whose leaves above the snow in winter have transiently reduced  $F_v/F_m$  compared with those that are covered by snow (Neuner et al. 1999). These findings suggest that *V. vitis-idaea* could benefit from the expected warmer winters and earlier snow melt in Finland, and much of the rest of its range (Rasmus et al. 2004 Henry 2008; Jylhä et al. 2008), by dehardening earlier and contributing more in the future to photosynthetic carbon gain in the understorey (Gehrmann et al., 2018; Larsen et al. 2007). However, this does not exclude the possibility that environmental changes involving reduced duration or size of snowpack can also be detrimental to understorey plants. Further north than our study, across a range of habitats in the Arctic where winters are colder than in central Finland, a reduced snow pack can cause sufficient cold stress to damage the leaves and shoots of shrubs including *V. vitis-idaea* (Preece and Phoenix 2013). Reduced snow cover can also lead to colder soils and prolonged freezing temperatures which can be a problem over a wider geographical range across Europe causing root mortality and affecting biogeochemical cycling (Groffman et al. 2001; Helama et al. 2011).

The link between increasing temperature and irradiance, and  $F_v/F_m$  recovery in *V. vitis-idaea* during dehardening reinforces established knowledge of this and other species (Ensminger et al. 2004; Havaux and Klopstech 2001; Muller et al. 2005; Neuner et al. 1999; Porcar-Castell et al. 2008;

Saarinen et al. 2016). At our site in central Finland, winter temperatures are relatively mild ( $-7.37 \pm 1.15$  °C; 10-year mean  $\pm 1$  SD January temperature, 2008-2018), and are expected to increase by 4-11 °C by the end of the century (Jylhä et al. 2008; Finland's Seventh National Communication under the United Nations Framework Convention on Climate Change 2017). Winter warming is forecast to increase productivity at the stand level (Porcar-Castell et al. 2011). However, the expected 30-40% reduction in snow cover (Jylhä et al. 2008; Finland's Seventh National Communication under the United Nations Framework Convention on Climate Change 2017) may lead to increased soil freezing or freeze thaw cycles which may partially negate the increase in productivity (Taulavuori et al. 2011). New estimates of ecosystem carbon assimilation aim to include the contribution of understorey vegetation in addition to that by tree species (Nikopainsius et al. 2015, Starr and Oberbauer, 2003). To estimate understorey photosynthesis during spring a better understanding of how it is affected by uneven snow cover under patchy canopies will be needed (Blume-Werry et al. 2016; Kreyling et al. 2012).

## 5. Conclusions (preferably 10-15 lines & no references)

In our study, there was a positive correlative relationship during the winter and spring between UV-screening by epidermal flavonols and  $F_v/F_m$  in leaves that were largely above the snowpack. More flavonols and anthocyanins accumulated in these upper-hummock leaves compared to leaves below the snowpack. The cross-protection provided by epidermal flavonols to cold and high light is likely to be regulated by solar radiation and temperature, but further research is needed to disentangle the roles of these two cues. Recent evidence that leaves can rapidly adjust their UV-screening to acclimate to light conditions (Barnes et al. 2017) suggests that a high-degree of fine tuning of these responses is possible, and our finding that flavonoids (flavonols and anthocyanins) accumulate in the upper hummocks even in cold conditions when metabolic activity is low supports this assertion. Leaves with a higher index of epidermal flavonols tended to have a lower epidermal anthocyanin index, and a compositional shift in this ratio towards flavonols during spring suggests fine-tuning of acclimation in response to increasing irradiance and a less cold

temperatures. Finally, on hummocks at our site in central Finland, leaves that remained under snow cover longer surpassed exposed leaves in their  $F_v/F_m$  directly following snowmelt despite having lower epidermal flavonols and anthocyanins but all leaves eventually reached a similar maximum photosynthetic capacity.

#### Funding

This study was funded by the Academy of Finland (grant number # 304519) to TMR. TS received grants from Finnish National Agency for Education (EDUFI fellowship program) and the Tiina and Herlin Foundation (project number 20180642).

#### Acknowledgments

Craig C Brelsford and Marieke Trasser helped to do the field work. We thank Timo Saarinen and Stef Bokhorst, for making their raw data available for our comparisons of  $F_v/F_m$ , light and temperature.

#### References

- Adams WW, Zarter CR, Ebbert V, Demmig-Adams B. 2004. Photoprotective Strategies of Overwintering Evergreens, *BioScience*. 54: 41–49.
- Agati G, Brunetti C, Di Ferdinando M, Ferrini F, Pollastri S, Tattini M. 2013. Functional roles of flavonoids in photoprotection: New evidence, lessons from the past. *Plant Physiology and Biochemistry* 72: 35–45.
- Barnes PW, Robson TM, Tobler MA, Bottger IN, Flint SD. 2017. Plant responses to fluctuating UV environments. Chapter 6 in B. Jordan. *Plant UV Biology*, CABI publishers 72–89.
- Barthod S, Cerovic Z, Epron D. 2007. Can dual chlorophyll fluorescence excitation be used to assess the variation in the content of UV-absorbing phenolic compounds in leaves of temperate tree species along a light gradient? *Journal of Experimental Botany*, 58: 1753–1760.
- Bilger W, Rolland M, Nybakken L. 2007. UV screening in higher plants induced by low temperature in the absence of UV-B radiation. *Photochem. Photobiol. Sci.* 6: 190–195.
- Blume-Werry G, Kreyling J, Laudon H, Milbau A. 2016. Short-term climate change manipulation effects do not scale up to long-term legacies: effects of an absent snow cover on boreal forest plants. *J. Ecol.* 104: 1638–1648.
- Bokhorst S, Bjerke JW, Bowles FW, Melillo J, Callaghan TV, Phoenix GK. 2008. Impacts of extreme winter warming in the sub-Arctic: growing season responses of dwarf shrub heathland. *Global Change Biology*. 14: 2603–2612.
- Bokhorst S, Bjerke JW, Davey MP, Taulavuori K, Taulavuori E, Laine K, Callaghan TV, and Phoenix GK. 2010. Impacts of extreme winter warming events on plant physiology in a sub-Arctic heath community. *Physiologia Plantarum*. 140: 128–140.



- 564 Brelsford CC, Morales LO, Nezval J, Kotilainen T, Hartikainen SM, Aphalo PJ, Robson TM. 2018. Do  
565 UV-A radiation and blue light during growth prime leaves to cope with acute high-light in  
566 photoreceptor mutants of *Arabidopsis thaliana*? *Physiol. Plant.* doi:10.1111/ppl.12749
- 567 Caldwell MM, Robberecht R, Flint SD. 1983. Internal filters: prospects for UV-acclimation in higher  
568 plants. *Physiol. Plant.* 58: 445–450.
- 569 Cerovic ZG, Masdoumier G, Ben Ghazlen M, Latouche G. 2012. A new optical leaf-clip meter for  
570 simultaneous non-destructive assessment of leaf chlorophyll and epidermal flavonoids, *Physiologia*  
571 *Plantarum* 146: 251–260.
- 572 Chalker-Scott L, James D Scott. 2004. Elevated Ultraviolet-B Radiation Induces Cross-protection to  
573 Cold in Leaves of *Rhododendron* under Field Conditions. *Photochemistry and Photobiology.* 79:  
574 199-204.
- 575 Coffey A, Prinsen E, Jansen MAK, and Conway J. 2017. The UVB photoreceptor UVR8 mediates  
576 accumulation of UV-absorbing pigments, but not changes in plant morphology, under outdoor  
577 conditions. *Plant, Cell Environment.* 40: 2250–2260.
- 578 Curl Jr. H, Hardy JT, Ellermeier R. 1972. Spectral absorption of Solar Radiation in Alpine Snow fields.  
579 *Ecology.* 53: 1189-1194.
- 580 Derksen C, Brown R, Mudryk L, and Luoju K. 2016. Terrestrial Snow (Arctic). In *State of the Climate*  
581 *in 2015. Bull. Am. Met. Soc.* 97: S145-S147.
- 582 Ensminger I, Sveshnikov D, Campbell DA, Funk C, Jansson S, Lloyd J, Shibistova O, Öquist G. 2004.  
583 Intermittent low temperatures constrain spring recovery of photosynthesis in boreal Scots pine  
584 forests. *Global Change Biology* 10: 995–1008.
- 585 Essery R, Pomeroy J, Ellis C and Link T. 2008. Modelling longwave radiation to snow beneath forest  
586 canopies using hemispherical photography or linear regression. *Hydrological Processes* 22: 2788–  
587 2800.
- 588 Gehrmann F, Hänninen H, Liu C Saarinen T. 2018. Phenological responses to small-scale spatial  
589 variation in snowmelt timing reveal compensatory and conservative strategies in subarctic-alpine  
590 plants, *Plant Ecology Diversity*, DOI: 10.1080/17550874.2018.1428693
- 591 Gorton, H. L. and Vogelmann, T. C. 2003. Ultraviolet Radiation and the Snow Alga *Chlamydomonas*  
592 *nivalis* (Bauer) Wille. *Photochemistry and Photobiology*, 77: 608-615.
- 593 Groffman PM, Driscoll CT, Fahey TJ, Hardy JP, Fitzhugh RD, Tierney GL. 2001. Colder soils in a warmer  
594 world: a snow manipulation study in a northern hardwood forest ecosystem. *Biogeochemistry.* 56:  
595 135– 150.
- 596 Harsch MA, HilleRisLambers J. 2016. Climate Warming and Seasonal Precipitation Change Interact to  
597 Limit Species Distribution Shifts across Western North America. *PLoS ONE* 11: e0159184.
- 598 Hartikainen SM, Jach A, Grané A. Robson TM. 2018 Assessing scale-wise similarity of curves with a  
599 thick pen: as illustrated through comparisons of spectral irradiance. *Ecology and Evolution in press*  
600 <https://doi.org/10.1002/ece3.4496>
- 601 Havaux M, Kloppstech K. 2001 The protective functions of carotenoid and flavonoid pigments  
602 against excess visible radiation at chilling temperature investigated in *Arabidopsis npq* and *tt*  
603 mutants. *Planta.* 213: 953–966
- 604 Hedstrom NR, Pomeroy JW. 1998. Measurements and modelling of snow interception in the boreal  
605 forest. *Hydrological processes* 12: 1611–1625.

- 606 Helama S, Tuomenvirta H, Venäläinen A. 2011. Boreal and subarctic soils under climatic change.  
607 Global and Planetary Change 79: 37–47
- 608 Henry HA. 2008. Climate change and soil freezing dynamics: historical trends and projected changes.  
609 Clim Change 87:421–434.
- 610 Hideg E, Jansen MA, Strid A. 2013. UV-B exposure, ROS, and stress: inseparable companions or  
611 loosely linked associates? Trends in Plant Science 18:107-115.
- 612 IPCC. 2014. Climate Change 2014. Impacts, Adaptation, and Vulnerability. Part A: Global and Sectoral  
613 Aspects. Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental  
614 Panel on Climate Change [Field, C.B. V.R. Barros, D.J. Dokken, K.J. Mach, M.D. Mastrandrea, T.E.  
615 Bilir, M. Chatterjee, K.L. Ebi, Y.O. Estrada, R.C. Genova, B. Girma, E.S. Kissel, A.N. Levy, S.  
616 MacCracken, P.R. Mastrandrea, and L.L. White (eds.)]. Cambridge University Press, Cambridge,  
617 United Kingdom and New York, NY, USA.
- 618 Ivanov AG, Sana PV, Hurry V, Öquist G, Huner NPA. 2008. Photosystem II reaction centre quenching:  
619 mechanisms and physiological role. Photosynth Res 98: 565–574.
- 620 Jansen MAK, Robson TM, Klem K, Urban O. 2017. UV-B induced morphological changes; an enigma.  
621 Chapter 5 in B Jordan. Plant UV Biology, CABI publishers, 58-71.
- 622 Johansson U, Gehrke C, Björn LO, Callaghan TV, Sonesson M. 1995. The effects of enhanced UV-B  
623 radiation on a sub-arctic heath ecosystem. Ambio 24 106-111.
- 624 Jylhä K, Fronzek S, Tuomenvirta H. et al. 2008. Changes in frost, snow and Baltic sea ice by the end of  
625 the twenty-first century based on climate model projections for Europe. Climate change. 86: 441.
- 626 Karlsson PS. 1985. Photosynthetic Characteristics and Leaf Carbon Economy of a Deciduous and an  
627 Evergreen Dwarf Shrub: *Vaccinium uliginosum* L. and *V. vitis-idaea* L. Holarctic Ecology 8: 9-17.
- 628 Kolari P, Peltoniemi M, Aurela, M, Böttcher, K, Tanis, Cemal Melih, Arslan, Ali Nadir. 2017.  
629 Phenological time lapse images from ground camera MC107 in Hyytiälä Pine stand (Version 2014-  
630 2017) [Data set]. Zenodo. <http://doi.org/10.5281/zenodo.1149217>
- 631 Kolb CA, Pfündel EE. 2005. Origins of non-linear and dissimilar relationships between epidermal UV  
632 absorbance and UV absorbance of extracted phenolics in leaves of grapevine and barley. Plant, Cell  
633 and Environment 25: 580–590.
- 634 Kreyling J. 2010. Winter climate change: a critical factor for temperate vegetation performance.  
635 Ecology 91:1939–1948.
- 636 Landi M, Tattini M, Gould KS. 2015. Multiple functional roles of anthocyanins in plant-environment  
637 interactions. Environmental and Experimental Botany, 19: 4-17.
- 638 Larsen K S, Ibrom A, Jonasson S, Michelsen A. Beier C. 2007. Significance of cold-season respiration  
639 and photosynthesis in a subarctic heath ecosystem in Northern Sweden. Global Change Biology,  
640 13: 1498-1508. doi:10.1111/j.1365-2486.2007.01370.x
- 641 Loik ME, Still CJ, Huxman TE. and Harte J. 2004. In situ photosynthetic freezing tolerance for plants  
642 exposed to a global warming manipulation in the Rocky Mountains, Colorado, USA. New  
643 Phytologist. 162: 331–341.
- 644 Lundell R, Saarinen T, Åström H, Hänninen H. 2008. The boreal dwarf shrub *Vaccinium vitis-idaea*  
645 retains its capacity for photosynthesis through the winter. Botany 86: 491–500.

- 646 Lütz C, Schönauer E, Neuner G. 2005. Physiological Adaptation Before and After Snow Melt in Green  
647 Overwintering Leaves of Some Alpine Plants. *Phyton*. 45: 139-156.
- 648 Mellander PE, Löfvenius MO, Laudon H. 2007. Climate change impact on snow and soil temperature  
649 in boreal Scots pine stands. *Climatic Change* 85: 179.
- 650 Morales LO, Tegelberg R, Brosché M, Lindfors A, Siipola S, Aphalo PJ. 2011. Temporal variation in  
651 epidermal flavonoids due to altered solar UV radiation is moderated by the leaf position in *Betula*  
652 *pendula*. *Physiologia Plantarum* 143: 261–270.
- 653 Muller O, Hikosaka K, Hirose T. 2005. Seasonal changes in light and temperature affect the balance  
654 between light utilisation and light harvesting components of photosynthesis in an evergreen  
655 understorey shrub. *Oecologia* 143: 501.
- 656 Murchie EH, Lawson T. 2013. Chlorophyll fluorescence analysis: a guide to good practice and  
657 understanding some new applications. *Journal of Experimental Botany*. 64: 3983–3998.
- 658 Neuner G, Ambach D, Aichner K. 1999. Impact of snow cover on photoinhibition and winter  
659 desiccation in evergreen *Rhododendron ferrugineum* leaves during subalpine winter. *Tree Physiol.*  
660 19:725-732.
- 661 Newsham KK, Robinson SA. 2009. Responses of plants in polar regions to UVB exposure: a meta-  
662 analysis. *Global Change Biology*. 15: 2574–2589.
- 663 Nikopensius M, Pisek J, Raabe K. 2015. Spectral reflectance patterns and seasonal dynamics of  
664 common understory types in three mature hemi-boreal forests. *International Journal of Applied*  
665 *Earth Observation and Geoinformation* 43: 84–91.
- 666 Nybakken L, Aubert S, Bilger W. 2004 Epidermal UV-screening of arctic and alpine plants along a  
667 latitudinal gradient in Europe. *Polar Biology*. 27: 391–398.
- 668 Perovich DK, 2007. Light reflection and transmission by a temperate snow cover. *Journal of*  
669 *Glaciology* 53: 201-211.
- 670 Petridis A, Döll S, Nichelmann L, Bilger W. and Mock H. 2016. *Arabidopsis thaliana* G2-LIKE  
671 FLAVONOID REGULATOR and BRASSINOSTEROID ENHANCED EXPRESSION1 are low-temperature  
672 regulators of flavonoid accumulation. *New Phytol.* 211: 912-925.
- 673 Phoenix GK, Gwynn-Jones D, Callaghan T, Sleep D. and Lee JA. 2001. Effects of global change on a  
674 sub-Arctic heath: effects of enhanced UV-B radiation and increased summer precipitation. *Journal*  
675 *of Ecology*, 89: 256-267.
- 676 Pomeroy JW, Gray DM, Hedström NR, Janowicz JR. 2002. Predictions of seasonal snow accumulation  
677 in cold climate forests. *Hydrological processes* 16: 3543–3558.
- 678 Porcar-Castell A, Juurola E, Nikinmaa E, Berninger F, Ensminger I Hari P. 2008. Seasonal acclimation  
679 of photosystem II in *Pinus sylvestris*. I. Using the rate constants of sustained thermal energy  
680 dissipation and photochemistry to study the effect of the light environment. *Tree physiology* 28:  
681 1475–1482.
- 682 Porcar-Castell A. 2011. A high-resolution portrait of the annual dynamics of photochemical and non-  
683 photochemical quenching in needles of *Pinus sylvestris*. *Physiologia Plantarum*. 143: 139-153.
- 684 Preece C, Phoenix GK. 2013. Impact of early and late winter icing events on sub-arctic dwarf shrubs.  
685 *Plant Biol.* 16: 125–132.

- 686 Rasmus S, Räisänen J, Lehning M. 2004. Estimating snow conditions in Finland in the late 21st  
687 century using the SNOWPACK model with regional climate scenario data as input. *Ann Glaciol* 38:  
688 238–244.
- 689 Richardson, S. Salisbury, F. 1977. Plant Responses to the Light Penetrating Snow. *Ecology*, 58: 1152-  
690 1158.
- 691 Robson TM, Hartikainen SM, Aphalo PJ. 2014. How does solar ultraviolet-B radiation improve  
692 drought tolerance of silver birch (*Betula pendula* Roth.) seedlings? *Plant Cell and Environment*, 38:  
693 953-67.
- 694 Saarinen T, Lundell R, Hänninen H. 2011. Recovery of photosynthetic capacity in *Vaccinium vitis-*  
695 *idaea* during mild spells in winter. *Plant Ecology*. 212: 1429-1440.
- 696 Saarinen T, Rasmus S, Lundell R, Kauppinen O-K, Hänninen H. 2016. Photosynthetic and phenological  
697 responses of dwarf shrubs to the depth and properties of snow. *Oikos*, 125: doi:10.1111/oik.02233
- 698 Salter WT, Turnbull TL, Rennenberg H, Adams MA. 2017. Research Solar UV upregulates  
699 photoprotection but slows photosynthesis in subalpine Australian plants. *Arctic, Antarctic, and*  
700 *Alpine*. 49: 673–685.
- 701 Schulz E, Tohge T, Zuther E, Fernie AR, Hinch DK. 2016. Flavonoids are determinants of freezing  
702 tolerance and cold acclimation in *Arabidopsis thaliana*. *Scientific Reports*. 6: 34027.
- 703 Searles PS, Flint SD, Caldwell MM. 2001. A meta-analysis of plant field studies simulating  
704 stratospheric ozone depletion. *Oecologia* 127: 1–10.
- 705 Semerdjieva SI, Pheonix GK, GK Hares, Gwynn-Jones D, Callaghan TV, Sheffield E. 2003a. Surface  
706 morphology, leaf and cuticle thickness of four dwarf shrubs from a sub-Arctic heath following long-  
707 term exposure to enhanced levels of UV-B. *Physiologia Plantarum*. 117: 289-294.
- 708 Semerdjieva SI, Sheffield E, Phoenix GK, Gwynn-Jones D, Callaghan, TV. Johnson GN. 2003b.  
709 Contrasting strategies for UV-B screening in sub-Arctic dwarf shrubs. *Plant, Cell Environment*, 26:  
710 957-964.
- 711 Siipola SM, Kotilainen T, Sipari N, Morales LO, Lindfers AV, Robson TM, Aphalo PJ. 2014. Epidermal  
712 UV-A absorbance and whole-leaf flavonoid composition in pea respond more to solar blue light  
713 than to solar UV radiation. *Plant Cell and Environment*, 38: 941–952.
- 714 Starr G, Oberbauer SF. 2003. Photosynthesis of arctic evergreens under snow: implications for  
715 tundra ecosystem carbon balance. *Ecology* 84: 1415–1420.
- 716 Sturm M, McFadden J, Liston GE, Chapin FS.III, Racine CH. Holmgren J. 2001. Snow-shrub  
717 interactions in Arctic tundra: A hypothesis with climatic implications. *Journal of Climate* 14: 336–  
718 344.
- 719 Sullivan JH, Gitz DC III, Liu-Gitz L, Xu C, Gao W, Slusser J. 2007. Coupling Short-Term Changes in  
720 Ambient UV-B levels with Induction of UV-Screening Compounds. *Photochemistry and*  
721 *Photobiology*, 83: 863–870.
- 722 Takahashi S, Badger MR. 2011. Photoprotection in plants: a new light on photosystem II damage.  
723 *Trends in Plant Science*. 16: 53-60.
- 724 Taulavuori K, Bauer E, Taulavuori E. 2011. Overwintering stress of *Vaccinium vitis-idaea* in the  
725 absence of snow cover. *Environmental and Experimental Botany* 72: 397– 403.

- 726 Taulavuori K, Hyöky V, Oksanen J, Taulavuori E, Julkunen-Tiitto R. 2016. Species-specific differences  
727 in synthesis of flavonoids and phenolic acids under increasing periods of enhanced blue light.  
728 *Environmental and Experimental Botany* 121: 145–150.
- 729 Finland's Seventh National Communication under the United Nations Framework Convention on  
730 Climate Change. 2017. Ministry of the Environment and Statistics Finland, Helsinki. 314 p.
- 731 Wang F. 2017. SIOX plugin in ImageJ: area measurement made easy. *UV4Plants Bulletin*, 2016(2), 37-  
732 44. <https://doi.org/https://doi.org/10.19232/uv4pb.2016.2.11>
- 733 Wargent JJ, Elfadly EM, Moore JP, Paul ND. 2011. Increased exposure to UV-B radiation during early  
734 development leads to enhanced photoprotection and improved long-term performance in *Lactuca*  
735 *sativa*. *Plant, Cell Environment*. 34: 1401–1413.
- 736 Öquist G, Huner NPA. 2003. Photosynthesis of Overwintering Evergreen Plants. *Annu. Rev. Plant Biol.*  
737 54: 329–55.
- 738

739 Tables

740 Table 1: Maximum and Minimum Air and Soil Temperatures, and maximum PAR (PPFD) on the days  
 741 of chlorophyll fluorescence and leaf pigments measurements. Freeze-thaw cycles are calculated for  
 742 the period between measurements (usually 2 weeks, including 14-02-2017) and represent the  
 743 number of times that the temperature passed above and returned below freezing during this period.  
 744 Tree trunk temperatures were recorded at breast height on the completed shaded north side of the  
 745 tree during winter and spring.

746 Table 2: Comparison of leaf pigment measurements and leaf morphological traits on both per area  
 747 and per mass basis. Final harvest of one-year old leaves (leaf age 394 days) and young leaves (leaf  
 748 age 29 days) leaves of *V.vitis-idaea* in July 2017. Mean  $\pm$  1 SE of 4 hummocks. Percentage changes of  
 749 upper and middle hummocks from lower hummock values are given in parentheses. ANOVA results  
 750 given in Table S5

751 List of Figures:  
 752 Graphical Abstract: Schematic representing the changes in leaf level processes moderated by the  
 753 snowpack, sunlight and temperature after different heights on hummocks of *V. vitis-idaea* during  
 754 spring dehardening.

755 Fig. 1: A. Time series of air and soil temperature with mean leaf temperature on dates of leaf trait  
 756 measurements. B. Daily time course of understorey photosynthetically active radiation (PAR) at 0.6  
 757 m height, at Hyytiälä Forestry Station; from the unfurling of leaves in June 2016 (followed until  
 758 November 2017), and including photosynthetic recovery period (June 2017 to November 2017) and  
 759 subsequent new leaf unfurling in June 2017.

760 Fig. 2: Solar spectral Irradiance at midday measured at the experimental site under a pine canopy in  
 761 a sunfleck and canopy shade, and in an adjacent open field, in clear sky conditions during the  
 762 measurement period on (A) 14-02-2017, (B) 07-03-2017 and (C) 19-06-2017. Spectral photon  
 763 irradiance is normalised to 1 at 780 nm (raw data at given in Fig. S2 and broken down in Table S1).

Fig. 3: Indices of (A) epidermal flavonols estimated from epidermal absorption at 375 nm; (B) epidermal anthocyanins, estimated from epidermal absorption at 515 nm, and (C) leaf chlorophyll estimated from the relative leaf transmission at 680 nm to 710 nm, on a leaf area basis. The period covered is from leaf emergence until leaves of *Vaccinium vitis-idaea* were one-and-a-half years old. Mean  $\pm$  1 SE of the 4 hummocks (n=4) are plotted for the adaxial (upper side) and abaxial (lower side) leaf epidermis, at the 3 heights on the hummock, and the period under the snowpack marked with vertical lines. \* indicates a significant difference with height, ANOVA summary Table S3.

Fig.4: (A) Time of course of the relationship between estimated epidermal flavonols and anthocyanins from Figs. 3A and B. Scatterplots and linear regression of (B) adaxial and (C) abaxial epidermal flavonols and anthocyanins, for all measurement dates together but the three hummock heights presented separately for comparison. \* indicates a significant difference with height, ANOVA summary Table S3.

Fig. 5: Time course of (A) maximum quantum yield of photosystem II ( $F_v/F_m$ ) and (B) effective quantum yield of photosystem II ( $\phi PS_{II}$ ) over the period from leaf emergence until leaves of *V. vitis-idaea* were 517 days old. Mean  $\pm$  1 SE of the 4 hummocks (n=4) are plotted, and the period under the snowpack marked with vertical lines. \* indicates a significant difference with height, ANOVA summary Table S3. The fluorescence parameters used for these calculations are given in Fig. S4.

Fig. 6: The relationship between  $F_v/F_m$  and (A) leaf temperature, (B) PAR and (C) epidermal flavonols at the time of measurement measured on six occasions. Each point is the mean of 10 measurements of each of 4 hummocks. These relationships only hold during the spring dehardening period (17-04-2017 to 18-05-2017) when the temperature is above freezing but leaves have not yet recovered to maximal  $F_v/F_m$  of 0.82.

787 Online Supplemental Material

788 Table S1: Spectral photon irradiance at the *V. vitis-idaea* hummocks. Spectral integrals and photo  
789 ratios calculated from spectral irradiance (Fig S2) measured in shade and sunflecks on three different  
790 dates (Feb, 14-02-2017; Mar 07-03-2017; Jun 19-06-17). The mean  $\pm$  1SE for the four hummocks are  
791 given plus the % decrease compared with adjacent open full sun measurements.

792 Table S2: Plant area index (PAI) and sunfleck duration on 20-07-2017 at the four hummocks sampled  
793 in the study, calculated from the hemispherical image analysis.

794 Table S3: Summaries of three and two-way ANOVA tables for the effects of hummock height, leaf  
795 side, leaf age, and their interactions on leaf pigments (Figs 3 and 4), chlorophyll fluorescence (Fig. 5),  
796 quenching (Figs S4 and S5), Leaf area and mass-based pigment values (Table 2) (df = degrees of  
797 freedom)

798 Table S4: Relationship between the maximum quantum yield of PSII ( $F_v/F_m$ ) and air and soil  
799 temperature tested for leaves at each hummock height. Linear regressions for leaves above and  
800 below the snowpack were tested against air and soil temperature respectively (2 weeks only). The  
801 mean of  $F_v/F_m$  over 4 hummocks on each of 6 measurement dates during the period of spring  
802 dehardening was used (leaf age 228-375, 24-03-2017 to 19-06-2017).

803 Table S5: Relationship between epidermal anthocyanins and flavonols for adaxial and abaxial  
804 epidermis separately. Each measurement date is tested separately for leaves at each hummock  
805 height.

806 Table S6: Relationships calculated from Saarinen et al, (2010) between air and soil temperatures and  
807  $F_v/F_m$  of *V. vitis-idaea* leaves at the time of measurement on date 2009-03-23.

808 Table S7: Relationships calculated from Bokhorst et al, (2010) between air and soil temperatures and  
809  $F_v/F_m$  of *V. vitis-idaea* leaves at the time of measurement in 2008 and 2009.



Figure S1: Grey scaled images and binary images with/without Floyd-Steinberg dithering created with IrfanView 4.44 for all the original hemispherical photos on the hummocks sampled at the measuring site on 20-07-2017.

Fig. S2: Solar spectral Irradiance at midday measured at the experimental site under a pine canopy in a sunfleck and canopy shade, and in an adjacent open field, in clear sky conditions during the measurement period on 14-02-2017, 07-03-2017 and 19-06-2017. (as Fig. 2 before normalisation). Presented according to date (left) and according to canopy position (right). Spectral integrals and photon ratios in Table S1.

Fig. S3: Ratio of adaxial to abaxial epidermal (A) flavonol and (B) anthocyanin indices, measured by Dualex. Ratios calculated from the data presented in Fig. 3, over the period from leaf emergence until leaves of *V. vitis-idaea* were one-and-a-half years old. Mean  $\pm$  1 SE of the 4 hummocks (n=4) are plotted at 3 hummock heights, and the period under the snowpack marked with vertical lines. \* indicates a significant difference with height on the adjacent date, ANOVA summaries in Table S3.

Fig. S4: The parameters  $F_m$  and  $F_0$  (A & B) are used to calculate  $F_v/F_m$ , the maximum quantum yield of photosystem II photochemistry (Fig 5A), and  $F_m'$  and  $F$  (C & D) are used to calculate  $\phi_{PSII}$  the effective quantum yield of photosystem II photochemistry (Fig 5B). The  $ETR$  (E), electron transfer rate is calculated from  $\phi_{PSII}$  and PAR incident on the leaf (F). The period covers leaf emergence until leaves of *V. vitis-idaea* were one-and-a-half years old is given. Mean  $\pm$  1 SE of the 4 hummocks (n=4) are plotted, and the period under the snowpack marked with vertical lines. Quenching coefficients calculated from these data in Fig. S5.

Fig. S4: A. The photochemical ( $qP$ ) and non-photochemical quenching (NPQ) and fraction of PSII centers that are 'open' ( $qL$ ) calculated from chlorophyll fluorescence measurements, defined according to Baker, 2008 (Chlorophyll Fluorescence: A Probe of Photosynthesis In Vivo, Annu. Rev. Plant Biol. 2008. 59:89–113 doi:10.1146/annurev.arplant.59.032607.092759). Time series over the

834 period from leaf emergence until leaves of *V. vitis-idaea* were one-and-a-half years old. Mean  $\pm$  1 SE  
835 of the 4 hummocks (n=4) are plotted, and the period under the snowpack marked with vertical lines.

Table 1: Maximum and Minimum Air and Soil Temperatures, and maximum PAR (PPFD) on the days of chlorophyll fluorescence and leaf pigments measurements. Freeze-thaw cycles are calculated for the period between measurements (usually 2 weeks, including 14-02-2017) and represent the number of times that the temperature passed above and returned below freezing during this period. Tree trunk temperatures were recorded at breast height on the completed shaded north side of the tree during winter and spring.

840  
841


Date of Measurement	DOY	Leaf Age Days	Max-Air Temperature °C	Min-Air Temperature °C	Max-Soil Temperature °C	Min-Soil Temperature °C	Max-PAR $\mu\text{mol m}^{-2}\text{s}^{-1}$	Freeze-thaw cycles	Snow Temperature °C	Tree Trunk Temperature (North)°C
14-02-2017	45	250	5.3	-1.3	0.5	0.3	112.9	34	--	--
07-03-2017	66	271	-0.4	-10.4	0.0	-1.2	289.5	8	-7.9	-6.6
24-03-2017	83	288	1.8	-4.4	0.4	0.2	474.7	14	-10.0	-4.2
06-04-2017	96	301	8.6	-0.3	0.5	0.4	764.4	15	-3.5	0.8
18-04-2017	108	313	3.7	-7.3	0.3	-0.2	677.0	--	-2.6	-1.4
19-04-2017	109	314	5.5	-5.9	0.4	-0.1	519.8	14	-2.7	2.2
02-05-2017	122	327	12.2	0.2	0.6	-0.4	828.9	--	-3.0	8.7
03-05-2017	123	328	10.9	0.6	0.8	0.5	868.8	17	-2.8	8.1
16-05-2017	136	341	13.5	-2.7	3.4	1.1	919.6	--	--	7.7
17-05-2017	137	342	10.4	1.6	3.7	2.1	271.1	16	--	6.6
30-05-2017	150	355	13.6	0.1	7.6	4.7	290.0	--	--	9.8
19-06-2017	170	375	20.9	9.7	11.9	9.9	932.3	3	--	--
08-07-2017	189	394/29	20.3	7.2	12.7	9.2	836.8	1	--	--
09-07-2017	190	395/30	21.9	9.9	13.9	10.3	899.6	--	--	--
08-11-2017	312	517/152	5.2	2.7	4.3	0.4	51.7	34	--	--

842

843 Table 2: Comparison of leaf pigment measurements and leaf morphological traits on per area and per mass basis. Final harvest of one-year old leaves (leaf  
 844 age 394 days) and young leaves (leaf age 29 days) leaves of *V.vitis-idaea* in July 2017. Mean of 4 hummocks  $\pm$  1 SE. Percentage changes of upper and  
 845 middle hummocks from lower hummock values are given in parentheses. ANOVA results given in Table S5.

Leaf Trait	Young leaves (Leaf Age 29 days old)			One-year old leaves (Leaf Age 394 days old)		
	Lower Hummock	Middle Hummock	Upper Hummock	Lower Hummock	Middle Hummock	Upper Hummock
<b>Morphological Trait</b>						
Leaf Area (mm <sup>2</sup> )	240 $\pm$ 9	240 $\pm$ 15 (0%)	246 $\pm$ 12.3 (+3%)	174 $\pm$ 11	206 $\pm$ 27 (+18%)	188 $\pm$ 20.6 (+8%)
Leaf Dry Matter Content (mg)	14.5 $\pm$ 1.9	13.6 $\pm$ 1.5 (-6%)	13.7 $\pm$ 1.6 (-5%)	28.9 $\pm$ 2.4	35.8 $\pm$ 4.8 (+24%)	34.9 $\pm$ 3.9 (+21%)
Leaf Mass Area (mg mm <sup>-2</sup> )	0.060 $\pm$ 0.006	0.056 $\pm$ 0.003 (-6%)	0.055 $\pm$ 0.005 (-8%)	0.166 $\pm$ 0.005	0.174 $\pm$ 0.01(+5%)	0.187 $\pm$ 0.01(+12%)
<b>Dualex (per area)</b>						
Chlorophyll index per leaf area (Dualex A mm <sup>-2</sup> )	78.6 $\pm$ 5.8	74.6 $\pm$ 6.4 (-5%)	77.8 $\pm$ 4.8 (-1%)	122.6 $\pm$ 5.6	136.8 $\pm$ 23.4 (+12%)	121.7 $\pm$ 17 (-1%)
Epidermal flavonol index per leaf area (Dualex A mm <sup>-2</sup> )	5.6 $\pm$ 0.7	6.8 $\pm$ 0.4 (+22%)	8.0 $\pm$ 0.7 (+43%)	5.7 $\pm$ 0.1	7.2 $\pm$ 0.9 (+25%)	6.8 $\pm$ 0.9 (+19%)
Epidermal anthocyanin index per leaf area (Dualex A mm <sup>-2</sup> )	0.89 $\pm$ 0.01	1.03 $\pm$ 0.07 (+16%)	1.07 $\pm$ 0.08 (+20%)	0.26 $\pm$ 0.03	0.36 $\pm$ 0.03 (+39%)	0.34 $\pm$ 0.02 (+29%)
<b>Dualex (per mass)</b>						
Chlorophyll index per leaf mass (Dualex A mg <sup>-2</sup> )	1346 $\pm$ 162	1322 $\pm$ 52 (-2%)	1445 $\pm$ 173 (+7%)	739 $\pm$ 14	798 $\pm$ 154 (+8%)	657 $\pm$ 99 (-11%)
Epidermal flavonol index per leaf mass (Dualex A mg <sup>-2</sup> )	93 $\pm$ 6	120 $\pm$ 4 (+30%)	146 $\pm$ 13 (+57%)	35 $\pm$ 1	42 $\pm$ 6 (+20%)	37 $\pm$ 5 (+6%)
Epidermal anthocyanin index per leaf mass (Dualex A mg <sup>-2</sup> )	15.2 $\pm$ 1.3	18.3 $\pm$ 0.6 (+20%)	19.4 $\pm$ 0.3 (+27%)	1.6 $\pm$ 0.2	2.1 $\pm$ 0.3 (+34%)	1.8 $\pm$ 0.1 (+15%)

Fig. 1: (A) Time series of air and soil temperature with mean leaf temperature on dates of leaf trait measurements. (B) Daily time course of understorey photosynthetically active radiation (PAR) at 0.6 m height, at Hyytiälä Forestry Station; from the unfurling of leaves in June 2016 (followed until November 2017), and including photosynthetic recovery period (June 2017 to November 2017) and subsequent new leaf unfurling in June 2017.



855 Fig. 2: Solar spectral Irradiance at midday measured at the experimental site under a pine canopy in  
 856 a sunfleck and canopy shade, and in an adjacent open field, in clear sky conditions during the  
 857 measurement period on (A) 14-02-2017, (B) 07-03-2017 and (C) 19-06-2017. Spectral photon  
 858 irradiance is normalised to 1 at 780 nm (raw data at given in Fig. S2 and broken down in Table S1).







Fig. 3: Indices of (A) epidermal flavonols estimated from epidermal absorption at 375 nm; (B) epidermal anthocyanins, estimated from epidermal absorption at 515 nm, and (C) leaf chlorophyll estimated from the relative leaf transmission at 680 nm to 710 nm, on a leaf area basis. The period covered is from leaf emergence until leaves of *Vaccinium vitis-idaea* were one-and-a-half years old. Mean  $\pm$  1 SE of the 4 hummocks (n=4) are plotted for the adaxial (upper side) and abaxial (lower side) leaf epidermis, at the 3 heights on the hummock, and the period under the snowpack marked with vertical lines. \* indicates a significant difference with height, ANOVA summary Table S3.



870



871



872

873



Fig.4: (A) Time of course of the relationship between estimated epidermal flavonols and anthocyanins from Figs. 3A and B. Scatterplots and linear regression of (B) adaxial and (C) abaxial epidermal flavonols and anthocyanins, for all measurement dates together but the three hummock heights presented separately for comparison. \* indicates a significant difference with height, ANOVA summary Table S3.







Fig. 5: Time course of (A) maximum quantum yield of photosystem II ( $F_v/F_m$ ) and (B) effective quantum yield of photosystem II ( $\phi_{PSII}$ ) over the period from leaf emergence until leaves of *V. vitis-idaea* were 517 days old. Mean  $\pm 1$  SE of the 4 hummocks (n=4) are plotted, and the period under the snowpack marked with vertical lines. \* indicates a significant difference with height, ANOVA summary Table S3. The fluorescence parameters used for these calculations are given in Fig. S4.




891 Fig. 6: The relationship between  $F_v/F_m$  and (A) leaf temperature, (B) PAR and (C) epidermal flavonols  
 892 at the time of measurement measured on six occasions. Each point is the mean of 10 measurements  
 893 of each of 4 hummocks. These relationships only hold during the spring dehardening period (17-04-  
 894 2017 to 18-05-2017) when the temperature is above freezing but leaves have not yet recovered to  
 895 maximal  $F_v/F_m$  of 0.82.



896



897



898